

Today's Plan:

Learning Target (standard): I will divide radical expressions by rationalizing the denominator and using the conjugate.

Students will: Complete practice problems over previous concepts at the boards, put up homework problems on the board and make necessary corrections to their own work, take notes over new material and complete practice problems over new concepts.

Teacher will: Provide practice problems over previous concepts, check homework problems for accuracy and provide students feedback, describe and provide examples of new concepts and assign students assessment problems over new concepts.

Assessment: Board work, homework check and homework assignment

Differentiation: Students will work at the board, go over and correct homework at their seats, actively engage in lecture over new concepts, practice new concepts with the aid of other students and the teacher and complete homework assignment.

Simplify:

$$\begin{aligned}
 \frac{4y}{\sqrt{6x^3}} \cdot \frac{\sqrt{6x^3}}{\sqrt{6x^3}} &= \frac{4y\sqrt{6x^3} \cancel{x \cdot x \cdot x}}{\sqrt{6 \cdot 6 \cdot x^3 \cdot x^3}} \\
 &= \frac{4xy\sqrt{6x}}{3\cancel{6}x^3} \\
 &= \frac{2y\sqrt{6x}}{3x^2}
 \end{aligned}$$

$\begin{array}{c} 6 \\ \sqrt{\quad} \\ 3 \cdot 2 \end{array}$

Simplify:

$$\frac{2}{\sqrt[5]{9b^2}} \cdot \frac{\sqrt[5]{3 \cdot 3 \cdot 3 \cdot b \cdot b \cdot b}}{\sqrt[5]{3 \cdot 3 \cdot 3 \cdot b \cdot b \cdot b}} = \frac{2 \sqrt[5]{27b^3}}{3b}$$

(Handwritten annotations: The denominator $\sqrt[5]{9b^2}$ is expanded to $\sqrt[5]{3 \cdot 3 \cdot b \cdot b}$. The numerator $\sqrt[5]{3 \cdot 3 \cdot 3 \cdot b \cdot b \cdot b}$ is expanded to $\sqrt[5]{3 \cdot 3 \cdot 3 \cdot b \cdot b \cdot b}$. Blue circles highlight the $3 \cdot 3$ in the denominator and the $3 \cdot 3 \cdot 3 \cdot b \cdot b \cdot b$ in the numerator, indicating cancellation. The final simplified expression is $\frac{2 \sqrt[5]{27b^3}}{3b}$.)