## Today's Plan:

**Learning Target (standard)**: I will find the derivative of trigonometric functions.

**Students will**: Complete practice problems over previous concepts at the boards, put up homework problems on the board and make neccessary corrections to their own work, take notes over new material and complete practice problems over new concepts.

**Teacher will**: Provide practice problems over previous concepts, check homework problems for accuarcy and provide students feedback, describe and provide examples of new concepts and assign students assessment problems over new concepts.

Assessment: Board work, homework check and homework assignment

**Differentiation**: Students will work at the board, go over and correct homework at their seats, actively engage in lecture over new concepts, practice new concepts with the aid of other students and the teacher and complete homework assignment.

p.386

$$40)y' = \frac{-2\sin x}{(\cos x + 1)^2}$$

$$35)y'' = 18\sec^4 3x + 36\sec^2 3x \tan^2 3x$$

$$36)y'' = 150\csc^4 5x \cot 5x + 150\csc^2 5x \cot^3 5x$$

$$41)y' = \frac{\sin y}{1 - x\cos y}$$

$$37)y'' = \sin x + x \cos x$$

$$42)y' = \frac{y}{-x}$$

$$39)y' = \frac{\sec^2 x}{2\sqrt{\tan x}}$$

44) 
$$y' = \frac{\sin(x-y) - y\sin x}{\sin(x-y) - \cos x}$$

Find the derivative:

$$f(x) = \cot 6x$$

$$f'(x) = -6csc^2 6x$$

Find the derivative:

$$y = \frac{\sec 2x}{\tan 2x + 1}$$

$$y' = (2 \sec 2x + \tan 2x)(\tan 2x + 1) - 2 \sec^2 2x (\sec 2x)$$
 $(\tan 2x + 1)^2$ 

$$= 2xc2x + tan^{2}2x + 2xc2x + tanx - 2xc^{3}2x$$

$$(+an2x+1)^{2}$$

$$y' = 2\sec 2x \left( + an^2 2x + + anx - 8c^2 2x \right)$$
 $\left( + an 2x + 1 \right)^2$ 

$$f(x) = (x^{6} + 1)^{5} (4x + 7)^{3}$$

$$f'(x) = 5(x^{6} + 1)^{4} (6x^{5})(4x + 7)^{3} + 3(4x + 7)^{2}(4)(x^{6} + 1)^{5}$$

$$= 30x^{5} (x^{6} + 1)^{4} (4x + 7)^{3} + 12(4x + 7)^{2} (x^{6} + 1)^{5}$$

$$= 6(x^{6} + 1)^{4} (4x + 7)^{2} [5x^{5} (4x + 7) + 2(x^{6} + 1)]$$

$$= 6(x^{6} + 1)^{4} (4x + 7)^{2} (20x^{6} + 35x^{5} + 2x^{6} + 2)$$

$$f'(x) = 6(x^{6} + 1)^{4} (4x + 7)^{2} (22x^{6} + 35x^{5} + 2)$$

Find the derivative:  

$$f(x) = \cos^4 x - 2x^2$$

$$f'(x) = 4\cos^3 x (-\sin x) - 4x$$

$$= -4\cos^3 x \sin x - 4x$$

$$f'(x) = -4(\cos^3 x \sin x + x)$$

## Find y'.

$$y^{4} + 3y - 4x^{3} = 5x + 1$$

$$4y^{3}y' + 3y' - 12x^{2} = 5$$

$$4y^{3}y' + 3y' = 12x^{2} + 5$$

$$y'(4y^{3} + 3) = 12x^{2} + 5$$

$$y'' = \frac{12x^{2} + 5}{4y^{3} + 3}$$

## Find the derivative:

$$f(x) = \frac{(3x^2 - \pi x)^4}{6} = \frac{1}{6} (3x^2 - \pi x)^4$$

$$f'(x) = \frac{2}{3} (3x^2 - \pi x)^3 (\log x - \pi)$$

$$f'(x) = \frac{2(3x^2 - \pi x)^3}{3} (\log x - \pi)$$

## Assignment:

Derivative Rules 2 #1-12